

Topological Solitons in Chern-Simons Theories for Double-Layer Fractional Quantum Hall Effect

Ikuo Ichinose¹ and Akira Sekiguchi

Institute of Physics, University of Tokyo, Komaba, Tokyo, 153 Japan

Abstract

Topological excitations in Chern-Simons (CS) gauge theories which describe double-layer fractional quantum Hall effect are studied. We shall consider generic (m, m, n) Halperin state. There are two type of solitons; one is vortex type excitation which has essentially the same structure with quasi-hole excitation in the single-layer case. The other is nontrivial pseudospin texture which is so-called skyrmion or meron. We shall first study qualitative properties of solitons in the original CS gauge theory and give results of numerical calculation. Then by using duality transformation, we derive an effective theory for topological excitations in the fractional quantum Hall effect. For spin texture, that theory is nonrelativistic CP^1 nonlinear σ -model with a CS gauge interaction and a Hopf term. Finally, we study quantum mechanical system of multi-soliton states, retaining only center coordinates of solitons as collective coordinates. Existence of inter-layer tunneling drastically changes excitation spectrum.

¹e-mail address: ikuo@hep1.c.u-tokyo.ac.jp

1 Introduction

Study of 2-dimensional electron system in a strong magnetic field is one of the most interesting topics in the condensed matter physics. Especially in the last few years, both experimentally and theoretically, existence of topologically stable excitations is suggested for quantum Hall state (QHS) with internal degrees of freedom like spin singlet QHS and QHS in double-layer systems. Theoretical study started with QHS at filling factor $\nu = 1$ by taking into account spin degrees of freedom[1]. It was suggested that low-energy charged excitation in that system is *not* a single-body electron but a topological spin texture with finite size and that excitation has similar structure with skyrmion in ferromagnet. At present, some experiments support this observation[2].

Somewhat detailed study on the double-layer case is given in Ref.[3] by using lowest Landau level (LLL) projection in the first-quantization language. Most of studies on double-layer QHS given so far are restricted to Halperin's (m, m, m) states.

In this paper, we shall study topological excitations in generic (m, m, n) states in the framework of the Chern-Simons (CS) gauge theory. In the CS gauge theory for QH effect, the original fermionic electron is expressed by bosonized electron with odd number of CS flux quanta attached[4]. QHS is characterized by the coherent condensation of bosonized electron. In the single layer case, there is a topological excitation, which has a close resemblance to the vortex in the superfluid, and that excitation is identified with Laughlin's quasi-hole excitation in the fractional quantum Hall (FQH) state[5]. In the QHS with an internal symmetry, other topological excitations are expected which have nontrivial structure in the internal space.

This paper is organized as follows. In Sec.2, we shall explain models, which describe FQH effect (FQHE) in double layer (DL) electron systems. In general, there are two CS gauge fields corresponding to $U(1) \otimes U(1)$ local symmetry, and two bosonized electron fields, which represent electrons in upper and lower layers, respectively. Ground state at specific filling factor has coherent condensation of both bosonized electron fields. In Sec.3, it is explained that there are essentially two types

of topological solitons in the models. One is vortex which has nontrivial winding number for $U(1)$ symmetry corresponding to the simultaneous phase rotation of the pseudospin up and down bosonized electron fields. The other is meron which has nontrivial topological structure in the $SU(2)$ internal space of pseudospin. Qualitative study of the topological solitons and numerical solutions to field equations are given. In Sec.4, an effective field theory for topological excitations is derived by using duality transformation. We first decompose bosonized electron fields into amplitude or density, common $U(1)$ phase factor and CP^1 variable. Lowest Landau level condition is obtained explicitly, which determines deviation of charge density in terms of topological currents of vortex and meron. Pseudospin CP^1 part of the effective field theory is a CP^1 nonlinear σ -model interacting with a CS gauge field and with a Hopf term. In Sec.5, quantum mechanics of solitons is discussed. Only center coordinates of them are taken into account as collective coordinates. Effect of inter-layer tunneling is considered. Single meron cannot exist with finite energy. Spectrum of meron pair is studied. Section 6 is devoted for discussion and conclusion.

2 Models

Lagrangian of the model is given as follows,

$$\begin{aligned} \mathcal{L}_\psi &= -\bar{\psi}_\uparrow(\partial_0 - ia_0^+ - ia_0^-)\psi_\uparrow - \bar{\psi}_\downarrow(\partial_0 - ia_0^+ + ia_0^-)\psi_\downarrow \\ &\quad - \frac{1}{2M} \sum_{\sigma=\uparrow,\downarrow} |D_j^\sigma \psi_\sigma|^2 - V[\psi_\sigma], \end{aligned} \quad (2.1)$$

$$\begin{aligned} \mathcal{L}_{\text{CS}} &= \mathcal{L}_{\text{CS}}(a_\mu^+) + \mathcal{L}_{\text{CS}}(a_\mu^-) \\ &= -\frac{i}{4}\epsilon_{\mu\nu\lambda} \left(\frac{1}{p} a_\mu^+ \partial_\nu a_\lambda^+ + \frac{1}{q} a_\mu^- \partial_\nu a_\lambda^- \right), \end{aligned} \quad (2.2)$$

where ψ_σ ($\sigma = 1, 2$ or $\uparrow, \downarrow$) is the bosonized electron fields in upper and lower layers, respectively, M is the mass of electrons and p and q are parameters. Greek indices take 0, 1 and 2, while roman indices take 1 and 2. $\epsilon_{\mu\nu\lambda}$ is the antisymmetric tensor

and the covariant derivative is defined as

$$D_j^{\uparrow\downarrow} = \partial_j - ia_j^+ \mp ia_j^- + ieA_j, \quad (2.3)$$

where external magnetic field is directed to the z -axis and in the symmetric Coulomb gauge $A_j = -\frac{B}{2}\epsilon_{jk}x_k$. $V[\psi_\sigma]$ is interaction term between electrons like the Coulomb repulsion and short-range four-body interaction, e.g.,

$$g_1(|\psi_\uparrow|^4 + |\psi_\downarrow|^4) + g_2|\psi_\uparrow\psi_\downarrow|^2, \text{ etc.} \quad (2.4)$$

Chern-Simons constraints are obtained by differentiating the Lagrangian with respect to $a_0^\pm$

$$\begin{aligned} \epsilon_{ij}\partial_i a_j^+ &= 2p\bar{\Psi}\Psi \\ \epsilon_{ij}\partial_i a_j^- &= 2q\bar{\Psi}\sigma_3\Psi \\ \Psi &= (\psi_\uparrow \ \psi_\downarrow)^t, \end{aligned} \quad (2.5)$$

where $\epsilon_{ij} = \epsilon_{0ij}$. As we shall see, $(p+q)$ must be an odd integer times π and $(p-q)$ must be an integer times π for the original electrons to be fermionic.

Partition function in the imaginary-time formalism is given as

$$Z = \int [D\bar{\psi} D\psi Da] \exp\left\{ \int d\tau d^2x (\mathcal{L}_\psi + \mathcal{L}_{\text{CS}}) \right\}. \quad (2.6)$$

The Lagrangian $\mathcal{L} = \mathcal{L}_\psi + \mathcal{L}_{\text{CS}}$ is invariant under $U(1) \otimes U(1)$ gauge transformation,

$$\begin{aligned} \Psi &\rightarrow e^{i\theta_1 + i\theta_2\sigma_3}\Psi \\ a_\mu^+ &\rightarrow a_\mu^+ + \partial_\mu\theta_1 \\ a_\mu^- &\rightarrow a_\mu^- + \partial_\mu\theta_2. \end{aligned} \quad (2.7)$$

Ground state for fractional quantum Hall effect is given by the following field configuration,

$$\psi_{\uparrow,0} = \sqrt{\bar{\rho}_\uparrow} = \sqrt{\bar{\rho}/2},$$

$$\psi_{\downarrow,0} = \sqrt{\bar{\rho}_{\downarrow}} = \sqrt{\bar{\rho}/2},$$

$$eB = \epsilon_{ij}\partial_i a_{j,0}^+ = 2p\bar{\rho}, \quad \epsilon_{ij}\partial_i a_{j,0}^- = 0, \quad a_{j,0}^+ = eA_j, \quad (2.8)$$

$$\nu = \frac{2\pi\bar{\rho}}{eB} = \frac{\pi}{p}, \quad (2.9)$$

where $\bar{\rho}$ is the average electron density. It is easily verified that the above static configuration is a solution to the field equations if and only if the filling factor ν has specific values given by (2.9).

One may think that there is a massless mode, because the configuration (2.8) breaks the $U(1) \otimes U(1)$ symmetry. However, these massless modes are eaten by the CS gauge fields and the CS gauge fields acquire mass term by the Higgs mechanism. This is important, because QHS must be incompressible and there must be no gapless modes.

It is not so difficult to show that the above ground state (2.8) corresponds to the following Halperin's (m, m, n) state[6] in the first-quantization language[7];

$$\begin{aligned} \Psi_G(z^\uparrow; z^\downarrow) &= \prod_{i<j} (z_i^\uparrow - z_j^\uparrow)^m \prod_{i<j} (z_i^\downarrow - z_j^\downarrow)^m \prod_{i,j} (z_i^\uparrow - z_j^\downarrow)^n \\ &\times \exp\left(-\frac{eB}{4}(\sum |z_i^\uparrow|^2 + \sum |z_i^\downarrow|^2)\right), \end{aligned} \quad (2.10)$$

where $m = \frac{1}{\pi}(p+q)$ and $n = \frac{1}{\pi}(p-q)$. For the specific case $m = n$, one can set $a_j^- = 0$ and only the single CS gauge field remains.

3 Solitons and numerical calculations

In this section, we shall study topological solitons in the CS gauge theory (2.1) and (2.2). We first assume the following specific form of the potential between bosonized electrons;

$$V[\psi_\sigma] = \frac{p}{M}(\bar{\Psi}\Psi)^2 + \frac{q}{M}(\bar{\Psi}\sigma_3\Psi)^2 + W[\psi_\sigma], \quad (3.1)$$

where $W[\psi_\sigma]$ represents long-range inter-layer and intra-layer Coulomb repulsions. It is argued that the above short-range repulsion is required in order to retain the

Pauli exclusion principle in the bosonization of fermionic field[8]. In the present case, field equation to be solved becomes very simple with this term as we shall see. It is expected, however, that the existence of this term influences behavior of solutions only near the center of them.

It is easily verified that the following modified Bogomol'nyi decomposition holds,

$$\begin{aligned} \sum_{\sigma} |D_j^{\sigma} \psi_{\sigma}|^2 &= \sum_{\sigma} |(D_1^{\sigma} - iD_2^{\sigma}) \psi_{\sigma}|^2 + \sum_{\sigma} \epsilon_{jk} \partial_j [\bar{\psi} i D_k^{\sigma} \psi_{\sigma}] \\ &\quad - \epsilon_{ij} \partial_i (a_j^+ - eA_j) (\bar{\Psi} \Psi) - \epsilon_{ij} \partial_i a_j^- (\bar{\Psi} \sigma_3 \Psi). \end{aligned} \quad (3.2)$$

By using the CS constraints (2.5) and (3.1), Hamiltonian (in the real-time formalism) is given as

$$\mathcal{H} = \frac{1}{2M} \sum_{\sigma} |(D_1^{\sigma} - iD_2^{\sigma}) \psi_{\sigma}|^2 + \frac{\omega_c}{2} (\bar{\Psi} \Psi) + W[\psi_{\sigma}], \quad (3.3)$$

where $\omega_c = eB/M$. If we treat the Coulomb interaction as a perturbation, the lowest-energy configurations satisfy the following “self-dual” equations,

$$(D_1^{\uparrow} - iD_2^{\uparrow}) \psi_{\uparrow} = 0, \quad (D_1^{\downarrow} - iD_2^{\downarrow}) \psi_{\downarrow} = 0. \quad (3.4)$$

Actually, (3.4) is nothing but the lowest Landau level condition, and there are many solutions to (3.4). This degeneracy is solved by the Coulomb interaction, and the ground state is the uniform configuration (2.8).

We are going to seek topological soliton solutions to the self-dual equation (3.4). In this section, we are interested in single soliton configurations. Therefore, we parameterize the fields as

$$\begin{aligned} \psi_{\uparrow} &= \sqrt{\bar{\rho}} \exp(u_1(r) + in_1 \theta(x)), \\ \psi_{\downarrow} &= \sqrt{\bar{\rho}} \exp(u_2(r) + in_2 \theta(x)), \end{aligned} \quad (3.5)$$

where $r = |\vec{x}|$, $\theta(x)$ is the azimuthal function $\theta(x) = \arctan\left(\frac{x_2}{x_1}\right)$, and n_1 and n_2 are integers which label topological solitons. By substituting (3.5) into (3.4), we have

$$\begin{aligned} a_k^+ + a_k^- - eA_k &= n_1 \partial_k \theta - \epsilon_{kl} \partial_l u_1, \\ a_k^+ - a_k^- - eA_k &= n_2 \partial_k \theta - \epsilon_{kl} \partial_l u_2. \end{aligned} \quad (3.6)$$

By differentiating Eq.(3.6) and using the CS constraint (2.5), we have the following Toda-type equations for $u_i(r)$ ($i = 1, 2$),

$$\frac{d^2 u_i}{d\lambda^2} + \frac{1}{\lambda} \frac{du_i}{d\lambda} + n_i \frac{\delta(\lambda)}{\lambda} + 2 - 2 \sum_j K_{ij} e^{2u_j} = 0, \quad (3.7)$$

$$\lambda = \sqrt{p\bar{\rho}} \cdot r = \frac{r}{\sqrt{2l_0}}, \quad K = \frac{1}{p} \begin{pmatrix} p+q & p-q \\ p-q & p+q \end{pmatrix}, \quad (3.8)$$

where l_0 is the magnetic length $l_0 = 1/\sqrt{eB}$. It is obvious that $u_1(r)$ and $u_2(r)$ couple with each other nontrivially unless $p = q$.

It is useful to rewrite (3.7) by introducing $v_i(\lambda) \equiv u_i(\lambda) + n_i \ln \lambda$ as follows;

$$\left(\frac{d^2}{d\lambda^2} + \frac{1}{\lambda} \frac{d}{d\lambda} \right) v_i + 2 - 2 \sum_j K_{ij} e^{2v_j} \lambda^{-2n_j} = 0, \quad (3.9)$$

where we have used the identity

$$\left(\frac{d^2}{d\lambda^2} + \frac{1}{\lambda} \frac{d}{d\lambda} \right) \ln \lambda = \frac{\delta(\lambda)}{\lambda}. \quad (3.10)$$

In terms of $v_i(\lambda)$,

$$\begin{aligned} \psi_{\uparrow} &= \sqrt{\bar{\rho}} \exp \left(v_1(\lambda) - n_1 \ln \lambda + i n_1 \theta(x) \right), \\ \psi_{\downarrow} &= \sqrt{\bar{\rho}} \exp \left(v_2(\lambda) - n_2 \ln \lambda + i n_2 \theta(x) \right). \end{aligned} \quad (3.11)$$

From (3.11), it is obvious that the integers n_i must be nonpositive, $n_i = 0, -1, -2, \dots$, for single-valuedness and regularity of solution at the center of soliton, and $v_i(\lambda)$'s are regular functions. On the other hand, boundary condition at the infinity is given as

$$\lim_{\lambda \rightarrow \infty} e^{u_i(\lambda)} = \frac{1}{\sqrt{2}}, \quad (3.12)$$

which guarantees that at sufficiently far from the soliton, amplitude of solution is equal to that of the ground state.

Before going into detailed study of the numerical calculation, let us examine qualitative properties of solitons. There are essentially two types of topological solitons.

The first one has a nontrivial winding number for $U(1)$ symmetry of simultaneous phase rotation of the both up and down bosonized electron fields. This soliton corresponds to the $(n_1, n_2) = (-1, -1)$ in (3.5), and has trivial structure for the pseudospin internal space. We shall call this soliton vortex. As we shall show in the numerical calculation, behavior of this soliton is very similar to the vortex in the single-layer system. The other ones are pseudospin texture and correspond to the $(-1, 0)$ and $(0, -1)$ in (3.5). Functions $u_1(\lambda)$ and $u_2(\lambda)$ for $(-1, 0)$ behave as

$$\begin{aligned} u_1(\lambda) &\sim \ln \lambda, \quad u_2(\lambda) \sim \text{const.}, \quad \text{for } \lambda \rightarrow 0, \\ u_1(\lambda) = u_2(\lambda) &\sim -\ln \sqrt{2}, \quad \text{for } \lambda \rightarrow \infty, \end{aligned} \quad (3.13)$$

and similarly for the $(0, -1)$. It is useful to introduce “pseudospin operator” $\vec{m}(x)$ as usual by¹,

$$\vec{m} = \bar{\Psi} \vec{\sigma} \Psi. \quad (3.14)$$

From (3.13), it is obvious for the $(-1, 0)$ meron

$$\begin{aligned} \vec{m}(x) &\sim -\hat{z}, \quad \text{as } \vec{x} \rightarrow 0, \\ \vec{m}(x) &\sim \cos \theta(x) \cdot \hat{x} + \sin \theta(x) \cdot \hat{y}, \quad \text{as } |\vec{x}| \rightarrow \infty, \end{aligned} \quad (3.15)$$

where $\hat{x}$, etc., are unit vectors in the x , y and z directions. Picture of the above configuration of $\vec{m}(x)$ is given in Fig.1. Similar picture is obtained for $(0, -1)$ soliton;

$$\begin{aligned} \vec{m}(x) &\sim \hat{z}, \quad \text{as } \vec{x} \rightarrow 0, \\ \vec{m}(x) &\sim \cos \theta(x) \cdot \hat{x} - \sin \theta(x) \cdot \hat{y}, \quad \text{as } |\vec{x}| \rightarrow \infty. \end{aligned} \quad (3.16)$$

It is obvious that they have fractional winding number in the $O(3)$ -pseudospin space, and we call them meron. A pair of $(-1, 0)$ and $(0, -1)$ merons form a skyrmion with unit topological number.²

¹ $\vec{m}(x)$ is not invariant under transformation (2.7). But under regular gauge transformation, topological structure of $\vec{m}(x)$ is invariant. Gauge-invariant definition of the topological charge will be given in Sec.4.

²More precisely, we should introduce pseudospin operator with unit length $\hat{m} \equiv \vec{m}/|\vec{m}|$. Then, topological charge density is given as $q(x) = \frac{1}{8\pi} \epsilon_{ij} \hat{m} \cdot [\partial_i \hat{m} \times \partial_j \hat{m}]$.

Let us introduce a couple of quantities; soliton charge Q is defined as

$$\begin{aligned}
Q &\equiv \int d^2x (\bar{\Psi}\Psi - \bar{\Psi}_0\Psi_0) \\
&= \int d^2x \frac{1}{2p} (\epsilon_{ij}\partial_i a_j^+ - \epsilon_{ij}\partial_i a_{j,0}^+) \\
&= \frac{1}{2p} \oint dx^j (a_j^+ - a_{j,0}^+) \\
&\equiv \frac{\Phi_{\text{CS}}}{2p},
\end{aligned} \tag{3.17}$$

where Ψ_0 etc., is the quantities of the ground state given by (2.8), and Φ_{CS} is the total CS flux of soliton. The electric charge Q_E is $Q_E = -eQ$. Similarly we define

$$\begin{aligned}
\bar{\Phi}_{\text{CS}} &\equiv \oint dx^j a_j^- \\
\bar{Q} &\equiv \frac{\bar{\Phi}_{\text{CS}}}{2q}.
\end{aligned} \tag{3.18}$$

Total angular momentum is given by

$$J = \int d^2x \epsilon_{ij} x_i \mathcal{P}_j, \tag{3.19}$$

$$\mathcal{P}_j = \frac{1}{2i} \left[\left\{ \bar{\psi}_\uparrow D_j^\dagger \psi_\uparrow - \bar{D}_j^\dagger \bar{\psi}_\uparrow \cdot \psi_\uparrow \right\} + \left\{ \bar{\psi}_\downarrow D_j^\dagger \psi_\downarrow - \bar{D}_j^\dagger \bar{\psi}_\downarrow \cdot \psi_\downarrow \right\} \right]. \tag{3.20}$$

It is useful to decompose the angular momentum as $J = U + V$,

$$\begin{aligned}
U &= \frac{1}{2i} \epsilon_{ij} \sum_\sigma \int d^2x x_i (\bar{\psi}_\sigma \partial_j \psi_\sigma - \partial_j \bar{\psi}_\sigma \psi_\sigma), \\
V &= -\epsilon_{ij} \int d^2x x_i (a_j^+ \bar{\Psi}\Psi - a_j^- \bar{\Psi}\sigma_3\Psi).
\end{aligned} \tag{3.21}$$

The CS constraint (2.5) is solved in the Coulomb gauge as follows,

$$\begin{aligned}
a_j^+(x) &= -\frac{p}{\pi} \epsilon_{jk} \int d^2y \frac{(x-y)^k}{|\vec{x}-\vec{y}|^2} \cdot \bar{\Psi}\Psi(y), \\
a_j^-(x) &= -\frac{q}{\pi} \epsilon_{jk} \int d^2y \frac{(x-y)^k}{|\vec{x}-\vec{y}|^2} \cdot \bar{\Psi}\sigma_3\Psi(y).
\end{aligned} \tag{3.22}$$

From (3.22),

$$V = -\frac{p}{2\pi} \left\{ \int d^2x \bar{\Psi}\Psi \right\}^2 - \frac{q}{2\pi} \left\{ \int d^2x \bar{\Psi}\sigma_3\Psi \right\}^2. \tag{3.23}$$

On the other hand for U , from (3.5) and (3.6)

$$U = \sum_{\sigma} n_{\sigma} \int d^2x |\psi_{\sigma}|^2. \quad (3.24)$$

Total angular momentum of soliton, $J(\text{soliton})$, is naturally defined as

$$\begin{aligned} J(\text{soliton}) &= J[\psi_{\sigma}] - J[\psi_{\sigma,0} = \sqrt{\bar{\rho}/2}] \\ &= \sum_{\sigma} n_{\sigma} \int d^2x |\psi_{\sigma}|^2 - \frac{p}{2\pi} \left[\left\{ \int d^2x \bar{\Psi} \Psi \right\}^2 - \left\{ \int d^2x \bar{\Psi}_0 \Psi_0 \right\}^2 \right] \\ &\quad - \frac{q}{2\pi} \left[\left\{ \int d^2x \bar{\Psi} \sigma_3 \Psi \right\}^2 - \left\{ \int d^2x \bar{\Psi}_0 \sigma_3 \Psi_0 \right\}^2 \right]. \end{aligned} \quad (3.25)$$

Similarly, energy of soliton is given as

$$\begin{aligned} E &= \int d^2x \left\{ \frac{1}{2M} \sum_{\sigma} |(D_1^{\sigma} - iD_2^{\sigma})\psi_{\sigma}|^2 + \frac{\omega_c}{2} \bar{\Psi} \Psi + W[\psi_{\sigma}] \right\} \\ &\quad - \int d^2x \left\{ \frac{\omega_c}{2} \bar{\Psi}_0 \Psi_0 + W[\psi_{\sigma,0}] \right\} \\ &= \int d^2x \left\{ W[\psi_{\sigma}] - W[\psi_{\sigma,0}] \right\}, \end{aligned} \quad (3.26)$$

where we have used the self-dual equation (3.4) and the fact that the total number of electrons is conserved. For general configuration $\hat{\psi}_{\sigma}(r)e^{in_{\sigma}\theta}$ of the winding numbers (n_1, n_2) and the CS gauge fields of the following form

$$a_k^{\pm} = \partial_k \theta \cdot a^{\pm}(r), \quad a_k^{\uparrow\downarrow} = a_k^{+} \pm a_k^{-} = \partial_k \theta \cdot a^{\uparrow\downarrow}(r), \quad (3.27)$$

the energy is explicitly given as

$$\begin{aligned} E &= 2\pi \int \frac{dr}{r} \left[\frac{1}{2M} \sum_{\sigma} \left\{ \left(r \frac{d\hat{\psi}_{\sigma}}{dr} \right)^2 - 2r(a^{\sigma} - \frac{eB}{2}r^2 - n_{\sigma})\hat{\psi}_{\sigma} \left(\frac{d\hat{\psi}_{\sigma}}{dr} \right) \right. \right. \\ &\quad \left. \left. + (a^{\sigma} - \frac{eB}{2}r^2 - n_{\sigma})^2 (\hat{\psi}_{\sigma})^2 \right\} + \frac{\omega_c}{2} r^2 (\bar{\Psi} \hat{\Psi} - \bar{\Psi}_0 \Psi_0) + r^2 (W[\hat{\psi}_{\sigma}] - W[\hat{\psi}_{\sigma,0}]) \right]. \end{aligned}$$

Therefore, for energy to be finite, the CS gauge fields must satisfy³

$$\begin{aligned} \lim_{r \rightarrow \infty} a^{+}(r) &= \frac{eB}{2}r^2 + \frac{1}{2}(n_1 + n_2), \\ \lim_{r \rightarrow \infty} a^{-}(r) &= \frac{1}{2}(n_1 - n_2). \end{aligned} \quad (3.28)$$

³For the (m, m, m) state, there is only single CS gauge field a_{μ}^{+} and $a_{\mu}^{-} = 0$. Then, merons have logarithmically divergent energy[3].

From (3.28),

$$Q = \frac{\pi}{p} \left(\frac{n_1 + n_2}{2} \right), \quad \bar{Q} = \int d^2x \left(\bar{\Psi} \sigma_3 \Psi - \bar{\Psi}_0 \sigma_3 \Psi_0 \right) = \frac{\pi}{q} \left(\frac{n_1 - n_2}{2} \right),$$

$$J(\text{soliton}) = \frac{1}{2} \left\{ \frac{\pi}{p} \left(\frac{n_1 + n_2}{2} \right)^2 + \frac{\pi}{q} \left(\frac{n_1 - n_2}{2} \right)^2 \right\}. \quad (3.29)$$

The above $J(\text{soliton})$ should be interpreted as spin of solitons, S . On the other hand, statistical parameter of soliton $\alpha(\text{soliton})$ is given as follows by the argument of the Aharonov-Bohm effect,

$$\alpha(\text{soliton}) = \alpha_+ + \alpha_-,$$

$$\alpha_+ = -\frac{Q\Phi_{\text{CS}}}{2} = -pQ^2,$$

$$\alpha_- = -\frac{\bar{Q}\bar{\Phi}_{\text{CS}}}{2} = -q\bar{Q}^2. \quad (3.30)$$

Therefore, the spin-statistics relation, $S = -\frac{1}{2\pi}\alpha$, holds generally for solitons in the present system. Actually, it is obvious from the above consideration that there are two CS gauge fields, a_μ^+ and a_μ^- , in the present model and they contribute to both spin and statistics *additively*. This result is rederived in later discussion in Sec.4 which is based on an effective theory of solitons.

Let us turn to the numerical solutions to the field equations (3.9). For example for the $(-1, -1)$ vortex, asymptotic solution near the center of soliton is obtained as

$$v_i(\lambda) \sim \beta_i - \frac{\lambda^2}{2}, \quad (3.31)$$

where β_i 's are parameters to be determined by the boundary condition at spatial infinity. Similar asymptotic solutions are obtained for general (n_1, n_2) . Using these asymptotic solutions, we obtain numerical solutions.

In the DL FQHE, there are more than one QHS's at the same filling factor. Actually as we showed in Sec.2, the present CS gauge theory describes QHS at $\nu = \pi/p$, which is independent of the parameter q . The value of q is determined by the inter and intra-layer Coulomb interactions $W[\psi_\sigma]$, which depend on inter-layer distance d . For each d , some specific Halperin state is a good variational wave function.

As an example, we shall consider the case $\nu = \frac{2}{5}$, and at this filling fraction, Halperin's states $(3, 3, 2)$, $(5, 5, 0)$ and $(1, 1, 4)$ are possible candidates for the QHS. From (2.10), it is obvious that the state $(5, 5, 0)$ corresponds to such a case that the inter-layer distance d is sufficiently large and the inter-layer Coulomb interaction is negligibly small. On the other hand for the $(1, 1, 4)$ state, the inter-layer Coulomb interaction dominates over the intra-layer interaction. Though this situation does not seem to be realized in samples in experiment so far, some interesting features of solitons appear in this QHS as we shall see. The state $(3, 3, 2)$ is in between.

Vortex solution ($n_1 = -1, n_2 = -1$) in the $(3, 3, 2)$ state is given in Fig.2. The picture shows that size of vortex is about four times of the magnetic length $l_0 = 1/\sqrt{eB}$, and its form is quite similar to that in the single-layer system[9]. Meron solutions ($n_1 = -1, n_2 = 0$) etc. are also given in Fig.2. It should be remarked that the density in the lower layer *increases* at the center of the meron. It is a result of the inter-layer Coulomb interaction. Size of the meron is about $4.5\sqrt{2}l_0$, and it is larger than the vortex. Creation energies of vortex and merons are calculation from these solutions and (3.26). We also show the numerical solutions of $(-2, 0)$ and $(-2, -1)$ solitons in Fig.2.

Similarly, meron solution for $(5, 5, 0)$ state is shown in Fig.3. As there is no inter-layer interaction, the density in the lower layer is not influenced by the deviation of density in the lower layer. Finally for the $(1, 1, 4)$ state, meron solution in Fig.4 has a long undulating tail because of the dominant inter-layer interaction.

From the above meron solutions, we can calculate normalized pseudospin vector $\hat{m}(x) = \vec{m}(x)/|\vec{m}(x)|$. Merons have a fractional topological charge in the $O(3)$ space. In the following section, we shall derive an effective field theory of solitons, and show which field theory describes $\hat{m}(x)$ and how the topological charge is related with electric charge of solitons. As we see, the lowest Landau level projection plays a very important role there.

Finally, we shall give wave functions corresponding to the topological solitons in

the first-quantization language. For soliton of (n_1, n_2) type which is located at $(w, \bar{w})$,

$$\begin{aligned}\Psi_S(z^\uparrow; z^\downarrow; w) &= \prod_{i < j} (z_i^\uparrow - z_j^\uparrow)^m \prod_{i < j} (z_i^\downarrow - z_j^\downarrow)^m \prod_{i, j} (z_i^\uparrow - z_j^\downarrow)^n \\ &\quad \times \prod_i (z_i^\uparrow - w)^{-n_1} \prod_i (z_i^\downarrow - w)^{-n_2} \\ &\quad \times \exp\left(-\frac{eB}{4}(\sum |z_i^\uparrow|^2 + \sum |z_i^\downarrow|^2)\right).\end{aligned}\quad (3.32)$$

4 Effective action of vortices and merons

In this section, we shall derive an effective field theory of topological solitons considered in Sec.3. At specific filling factor (2.9), these solitons are excitations of low energies. Lowest Landau level condition is expressed in terms of topological current of these solitons, as we shall see in later discussion.

We first parameterize the bosonized electron fields as follows[10],

$$\begin{aligned}\psi_\sigma &= J_0^{1/2} \phi z_\sigma, \\ \phi &= e^{i\tilde{\theta}} \phi_v \in U(1), \quad z = (z_1 \ z_2)^t \in CP^1,\end{aligned}\quad (4.1)$$

where $e^{i\tilde{\theta}}$ is a regular part and ϕ_v is a topologically nontrivial part, which represents vortex degrees of freedom. CP^1 variable z_σ represents pseudospin degrees of freedom and satisfies CP^1 condition $\sum |z_\sigma|^2 = 1$, and ψ_σ is invariant under $z_\sigma \rightarrow e^{i\alpha} z_\sigma$ and $\phi \rightarrow e^{-i\alpha} \phi$.

Substituting (4.1) into (2.1),

$$\begin{aligned}\mathcal{L}_\psi &= -J_0 [\bar{\phi} \partial_0 \phi + (\bar{z} \partial_0 z) - ia_0^+ - ia_0^- (\bar{z} \sigma_3 z)] \\ &\quad - \frac{J_0}{2M} [\bar{\phi} \partial_j \phi + \bar{z} \partial_j z - i(a_j^+ - eA_j) - ia_j^- (\bar{z} \sigma_3 z)] \\ &\quad \times [\partial_j \bar{\phi} \cdot \phi + \partial_j \bar{z} \cdot z + i(a_j^+ - eA_j) + ia_j^- (\bar{z} \sigma_3 z)] + \mathcal{L}_z, \\ \mathcal{L}_z &= -\frac{J_0}{2M} [\overline{D_j z} \cdot D_j z + (\bar{z} D_j z)^2],\end{aligned}\quad (4.2)$$

where

$$D_\mu z = (\partial_\mu - ia_\mu^- \sigma_3) z, \quad (4.3)$$

and we have neglected derivative terms of J_0 , for J_0 is *not* a dynamical variable⁴ and its value is determined by LLL condition as we shall see. We have also neglected the interaction term $V[\psi_\sigma]$, which will be discussed later on.

We shall perform duality transformation[11]. First, we use the following identity;

$$e^{-\int d\tau d^2x \frac{K}{2} \varphi_j \varphi_j} = \int [DJ] e^{-\int d\tau d^2x (\frac{1}{2K} J_j J_j + i J_j \varphi_j)}. \quad (4.4)$$

In the present case, $K = \frac{\bar{\rho}}{M}$ and

$$\varphi_j = \bar{\phi} \frac{\partial_j}{i} \phi + \bar{z} \cdot \frac{\partial_j}{i} z - (a_j^+ - e A_j) - a_j^- (\bar{z} \sigma_3 z), \quad (4.5)$$

and the Lagrangian $\mathcal{L}$ is expressed as

$$\begin{aligned} \mathcal{L} = & -J_0 [\bar{\phi} \partial_0 \phi + (\bar{z} \partial_0 z) - i a_0^+ - i a_0^- (\bar{z} \sigma_3 z)] - \frac{1}{2K} \vec{J} \cdot \vec{J} - i J_j [\bar{\phi} \frac{\partial_j}{i} \phi + \bar{z} \cdot \frac{\partial_j}{i} z] \\ & + i J_j (a_j^+ - e A_j) + i J_j a_j^- (\bar{z} \sigma_3 z) + \mathcal{L}_z + \mathcal{L}_{\text{CS}}. \end{aligned} \quad (4.6)$$

It is obvious that integration over $\tilde{\theta}$ gives the divergentless condition

$$\partial_\mu J_\mu = 0. \quad (4.7)$$

Solution to (4.7) is

$$J_\mu = \frac{1}{2\pi} \epsilon_{\mu\nu\lambda} \partial_\nu b_\lambda. \quad (4.8)$$

Substituting (4.8) into (4.6),

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2\pi} \epsilon_{ij} \partial_i b_j [\bar{\phi}_v \partial_0 \phi_v + (\bar{z} \partial_0 z) - i a_0^+ - i a_0^- (\bar{z} \sigma_3 z)] \\ & - \frac{1}{2K} \frac{1}{(2\pi)^2} (\partial_0 b_j - \partial_j b_0)^2 - \frac{i}{2\pi} \epsilon_{ij} (\partial_0 b_i - \partial_i b_0) [\bar{\phi}_v \frac{\partial_j}{i} \phi_v + \bar{z} \cdot \frac{\partial_j}{i} z] \\ & + \frac{i}{2\pi} \epsilon_{ij} (\partial_0 b_i - \partial_i b_0) (a_j^+ - e A_j + a_j^- (\bar{z} \sigma_3 z)) + \mathcal{L}_z + \mathcal{L}_{\text{CS}}. \end{aligned} \quad (4.9)$$

Next we integrate over b_0 with the Coulomb gauge $\partial_j b_j = 0$. The form of the integrand is the following,

$$\begin{aligned} & -\frac{1}{2K} \frac{1}{(2\pi)^2} (\partial_j b_0)^2 + i b_0 \Phi, \\ \Phi = & -\frac{1}{2\pi} \epsilon_{ij} \partial_i [\bar{\phi}_v \frac{\partial_j}{i} \phi_v + \bar{z} \cdot \frac{\partial_j}{i} z - (a_j^+ - e A_j) - a_j^- (\bar{z} \sigma_3 z)]. \end{aligned} \quad (4.10)$$

⁴It is obvious that the term $J_0^{1/2} \partial_0 J_0^{1/2}$ is total derivative

Integration gives

$$\int [Db_0] e^{\int d^3x [-C(\partial_j b_0)^2 + ib_0 \Phi]} = e^{-\frac{1}{4} \int \Phi \frac{1}{C} \Delta^{-1} \Phi}, \quad (4.11)$$

where $C = \frac{1}{2K(2\pi)^2}$ and

$$\int \Phi \Delta^{-1} \Phi = -\frac{1}{2\pi} \int d\tau \int d^2x d^2y \Phi(x) \ln |\vec{x} - \vec{y}| \Phi(y), \quad (4.12)$$

$$\Delta^{-1}(x) = \int \frac{d^2k}{(2\pi)^2} \frac{e^{ik \cdot x}}{k^2}. \quad (4.13)$$

We furthermore integrate over the CS fields a_0^+ which gives the CS constraint,

$$\epsilon_{ij} \partial_i a_j^+ = 2p |\Psi|^2 = 2p J_0 = \frac{p}{\pi} \epsilon_{ij} \partial_i b_j, \quad (4.14)$$

and Φ is expressed as

$$\Phi = -\frac{1}{2\pi} \epsilon_{ij} \partial_i \left[\bar{\phi}_v \frac{\partial_j}{i} \phi_v + \bar{z} \cdot \frac{D_j}{i} z - \left(\frac{p}{\pi} b_j - e A_j \right) \right]. \quad (4.15)$$

Let us consider quantization of the field b_j . Kinetic term is given as $(\partial_0 b_j)^2$ and a system is simply a harmonic oscillator with the following form of the action $L = \dot{x}^2 + (x - a)^2$. We shall consider static configuration of solitons. Then, Lagrangian of b_j field, $\mathcal{L}_b$, is given by

$$\mathcal{L}_b = -C(\partial_0 b_j)^2 - \frac{1}{4C} \int d^2y \Phi(x, \tau) \Delta^{-1}(x - y) \Phi(y, \tau). \quad (4.16)$$

It is convenient to take $B(x) \equiv \epsilon_{ij} \partial_i b_j$ as dynamical variable for quantization in stead of $b_j(x)$ itself which is subject to the Coulomb gauge condition. The original field b_j is expressed in terms of B as

$$b_i(x) = \epsilon_{ij} \partial_j^x \int d^2y \Delta^{-1}(x - y) B(y). \quad (4.17)$$

We shall introduce Fourier transformed variables as

$$\begin{aligned} b_i(x) &= \int \frac{d^2q}{(2\pi)^2} e^{iq \cdot x} \tilde{b}_i(q), \\ B(x) &= \int \frac{d^2q}{(2\pi)^2} e^{iq \cdot x} \tilde{B}(q), \text{ etc.} \end{aligned} \quad (4.18)$$

In terms of Fourier transformed fields,

$$\begin{aligned}\int d^2x (\partial_0 b_j)^2 &= \int \frac{d^2q}{(2\pi)^2} \frac{1}{q^2} \partial_0 \tilde{B}(q) \partial_0 \tilde{B}(-q), \\ \int d^2x d^2y \Phi(x) \Delta^{-1}(x-y) \Phi(y) &= \int \frac{d^2q}{(2\pi)^2} \tilde{\Phi}(q) \frac{1}{q^2} \tilde{\Phi}(-q).\end{aligned}\quad (4.19)$$

Then, returning to the real-time formalism as $t = -i\tau$, canonical conjugate variable of $\tilde{B}(q)$ is

$$\Pi_B(q) = C \frac{2}{(2\pi)^2 q^2} \partial_t \tilde{B}(-q). \quad (4.20)$$

Hamiltonian is given as

$$H_B = \int d^2q \left[\frac{K}{2} q^2 (2\pi)^4 \Pi_B(q) \Pi_B(-q) + \frac{K}{2} \tilde{\Phi}(q) \frac{1}{q^2} \tilde{\Phi}(-q) \right], \quad (4.21)$$

and

$$\tilde{\Phi}(q) = \frac{1}{2\pi} \frac{1}{\nu} \tilde{B}(q) + \dots \quad (4.22)$$

From (4.21) and (4.22), it is obvious that the ground state is the configuration which satisfies $\Phi = 0$ and excitation energies are given by $N\omega_c$, where N is the number of excitations and $\omega_c = eB/M$.

Ground state condition $\Phi = 0$ is nothing but the LLL condition, which is given as follows[1],

$$J_0^v + J_0^S - \frac{2p}{(2\pi)^2} \epsilon_{ij} \partial_i b_j + \frac{e}{2\pi} \epsilon_{ij} \partial_i A_j = 0, \quad (4.23)$$

where topological densities and currents are defined as

$$J_\mu^v = \frac{1}{2\pi} \epsilon_{\mu\nu\lambda} \partial_\nu \left(\bar{\phi}_v \frac{\partial_\lambda}{i} \phi_v \right), \quad (4.24)$$

$$J_\mu^S = \frac{1}{2\pi} \epsilon_{\mu\nu\lambda} \partial_\nu \left(\bar{z} \frac{D_\lambda}{i} z \right). \quad (4.25)$$

There is another $U(1) \otimes U(1)$ gauge-invariant topological current, which is defined as

$$J_\mu^3 = \frac{1}{2\pi} \epsilon_{\mu\nu\lambda} \partial_\nu \left(\bar{z} \sigma_3 \frac{\tilde{D}_\lambda}{i} z \right), \quad (4.26)$$

where $\tilde{D}_\lambda = \partial_\lambda - i(a_\lambda^+ - eA_\lambda)$. This topological charge distinguishes between $(-1, 0)$ and $(0, -1)$ merons.

In the small-size or the long-distance limit of the topological solitons, the above currents are given as

$$\begin{aligned}
J_0^v(x) &= \sum_{\alpha} q_{\alpha}^v \delta(x - x^{v,\alpha}), \\
J_j^v(x) &= \sum_{\alpha} q_{\alpha}^v \dot{x}_j^{v,\alpha} \delta(x - x^{v,\alpha}), \\
J_0^S(x) &= \sum_{\alpha} q_{\alpha}^S \delta(x - x^{S,\alpha}), \\
J_j^S(x) &= \sum_{\alpha} q_{\alpha}^S \dot{x}_j^{S,\alpha} \delta(x - x^{S,\alpha}),
\end{aligned} \tag{4.27}$$

where index α distinguishes solitons and q_{α}^v and q_{α}^S are topological charges of solitons, i.e.,

$$\begin{aligned}
q^v &= -1 \text{ for } (-1, -1) \text{ vortex,} \\
q^S &= -\frac{1}{2} \text{ for } (-1, 0), (0, -1) \text{ merons,} \\
q^S &= +\frac{1}{2} \text{ for } (1, 0), (0, 1) \text{ merons,}
\end{aligned} \tag{4.28}$$

and $x^{v,\alpha}$ and $x^{S,\alpha}$ are their coordinates. From (3.11), merons $(1, 0)$ etc. do not exist as a regular solution to the self-dual equation. This situation is similar to the single-layer system, in which quasi-particle excitation does not exist as a regular solution. However, these solitons must exist as low-energy excitations and their behavior away from the center is described by (3.11).

The LLL condition (4.23) determines the density of electrons in terms of the topological currents. For example from (4.14), the vortex $(-1, -1)$ has charge νe and the merons $(-1, 0)$ and $(0, -1)$ have charge $\frac{\nu e}{2}$.

Here, a few remarks about the LLL condition are in order. The condition (4.23) is correct for static configurations. For moving soliton, it is not so difficult to show that there appears additional term which is proportional to $\epsilon_{ij} \partial_i (J_j^v + J_j^S)$. Existence of this term, however, does not change the discussion on statistics of solitons, etc, given below. For example,

$$\int_{S_0} d^2x \epsilon_{ij} \partial_i J_j = \int_{\partial S_0} J_j dx_j, \tag{4.29}$$

where S_0 is a domain and ∂S_0 is its boundary. For sufficiently large domain whose boundary is far away from solitons, the above integral is vanishingly small and therefore charge of solitons is determined solely by the topological density.

There is obviously an ambiguity in the decomposition (4.1), i.e., the bosonized electron fields are invariant under the following transformation,

$$\phi \rightarrow e^{-i\varphi} \phi, \quad z_\sigma \rightarrow e^{i\varphi} z_\sigma, \quad (4.30)$$

where the phase φ can have a *nontrivial* winding number. This singular transformation is allowed at the center of solitons with topological numbers $n_1 \neq 0$ and $n_2 \neq 0$, as the amplitude J_0 vanishes there. From the definition of the topological charges (4.24) and (4.25),

$$\begin{aligned} J_0^v &\rightarrow J_0^v - \frac{1}{2\pi} \epsilon_{ij} \partial_i \partial_j \varphi, \\ J_0^S &\rightarrow J_0^S + \frac{1}{2\pi} \epsilon_{ij} \partial_i \partial_j \varphi. \end{aligned} \quad (4.31)$$

Therefore, the total topological charge is invariant under (4.30).

With the LLL condition (4.23), the Lagrangian is given as

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2\pi} \epsilon_{ij} \partial_i b_j \left[\bar{\phi}_v \partial_0 \phi_v + (\bar{z} \partial_0 z) - i a_0^- \cdot (\bar{z} \sigma_3 z) \right] - \frac{i}{2\pi} \epsilon_{ij} \partial_0 b_i \left[\bar{\phi}_v \frac{\partial_j}{i} \phi_v + \bar{z} \cdot \frac{\partial_j}{i} z \right] \\ & + \frac{i}{2\pi} \epsilon_{ij} \partial_0 b_i \cdot (a_j^+ - e A_j) + \frac{i}{2\pi} \epsilon_{ij} \partial_0 b_i a_j^- \cdot (\bar{z} \sigma_3 z) + \mathcal{L}_z + \mathcal{L}_{\text{CS}}(a_\mu^-). \end{aligned} \quad (4.32)$$

After some calculation, it is shown

$$\text{1st+2nd+4th terms of (4.32)} = -i b_i \cdot (J_i^v + J_i^S). \quad (4.33)$$

On the other hand, the third term is evaluated as follows;

$$\frac{i}{2\pi} \epsilon_{ij} \partial_0 b_i \cdot (a_j^+ - e A_j) \Rightarrow -\frac{i}{2\pi} \epsilon_{ij} b_i \cdot \partial_0 (a_j^+ - e A_j). \quad (4.34)$$

Let us denote $\hat{J}_\mu = J_\mu^v + J_\mu^S$. In the Coulomb gauge,

$$(a_j^+ - e A_j)(x) = 2\pi \int d^2 y \epsilon_{jk} \partial_k^x \Delta^{-1}(x - y) \hat{J}_0(y). \quad (4.35)$$

By the conservation of the topological currents

$$\partial_\mu \hat{J}_\mu = 0, \quad (4.36)$$

$$\begin{aligned} \epsilon_{ij} \partial_0 (a_j^+ - eA_j)(x) &= -2\pi \int d^2y \partial_i^x \Delta^{-1}(x-y) \partial_0 \hat{J}_0(y) \\ &= 2\pi \int d^2y \partial_i^x \partial_k^x \Delta^{-1}(x-y) \cdot \hat{J}_k(y). \end{aligned} \quad (4.37)$$

On the other hand,

$$b_i(x) = \frac{2\pi^2}{p} \int d^2y \epsilon_{ij} \partial_j^x \Delta^{-1}(x-y) \cdot \hat{J}_0(y) + \frac{\pi}{p} eA_i(x). \quad (4.38)$$

Then from (4.37) and (4.38)

$$\begin{aligned} \int d^2x b_i(x) \epsilon_{ij} \partial_0 (a_j^+ - eA_j)(x) &= \frac{2\pi^2}{p} \int d^2x d^2y d^2z \left[\partial_i^x \partial_k^x \Delta^{-1}(x-z) \cdot \hat{J}_k(z) \epsilon_{ij} \partial_j^x \Delta^{-1}(x-y) \right. \\ &\quad \left. \times \hat{J}_0(y) \right] + \frac{\pi}{p} \int d^2x d^2z eA_i(x) \partial_i^x \partial_k^x \Delta^{-1}(x-z) \cdot \hat{J}_k(z) \\ &= 0, \end{aligned} \quad (4.39)$$

where we have used the Coulomb gauge condition.

Therefore, the effective Lagrangian $\mathcal{L}_E$ has the following final form;

$$\mathcal{L}_E = \mathcal{L}_z + \mathcal{L}_{\text{CS}}(a_\mu^-) - ib_i \cdot \hat{J}_i. \quad (4.40)$$

With the LLL condition (4.23), the term (4.33) is nothing but the Hopf term which determines statistics of solitons[12]. For vortex with $(-1, -1)$, J_μ^v has charge -1 . It is not difficult to show that statistical parameter of that excitation coming from the above Hopf term is $\pi\nu$, where the filling factor ν is given by $\nu = \frac{2\pi\bar{\rho}}{eB} = \frac{\pi}{p}$. On the other hand for merons $(-1, 0)$ and $(0, -1)$, J_μ^S has charge $-1/2$, and then the statistical parameter from the Hopf term is $\pi\nu(\frac{1}{2})^2$. The above results are consistent with the discussion in terms of the Aharonov-Bohm effect in Sec.3, i.e., the Hopf term in (4.40) gives the statistical parameter α_+ in (3.30).

The above Hopf term also gives spin to solitons[12], which is consistent with the spin-statistics relation. As we showed in Sec.3, merons in the present system have

the extra terms both in their spin and statistical parameter, besides those coming from the Hopf term. These terms come from the coupling with the CS field a_μ^- , as we can see from (4.40) and the discussion in Sec.3. Actually recently, CS gauge theory, which is a relativistic version of (4.40) *without* Hopf term, is studied[13]. There, it is shown that solitons similar to those in the present system have a fractional spin and also fractional statistics. We also studied topological solitons in the effective theory (4.40) very recently[14] and found that the CS gauge field a_μ^- gives spin and statistical parameter to merons which are consistent with the results in Sec.3.

In the following section, we shall consider the quantization of multi-soliton configurations retaining only center coordinates of solitons. In that discussion, it will become clear that the above Hopf term surely determines statistics of solitons.

5 Quantum mechanics of topological excitations

In the previous section, we have obtained the effective Lagrangian of topological solitons (4.40) by the duality transformation. Also as we showed by the numerical calculation in Sec.3, topological solitons have a finite size of order of the magnetic length. Actually form of the meron is determined by the terms $\mathcal{L}_z + \mathcal{L}_{\text{CS}}(a_\mu^-)$ in $\mathcal{L}_E$, and long-wave-length properties of topological excitations are determined by the term $b_i \cdot \hat{J}_i$. Especially, by using the topological currents in the form of the small-size limit (4.27), one can easily obtain an effective Lagrangian of multi-soliton system from $b_i \cdot \hat{J}_i$, which retains only coordinates of solitons as collective coordinates.

The LLL condition is solved as (4.38). It is not so difficult to calculate the long-distance limit of the term $b_i \cdot \hat{J}_i$,

$$\begin{aligned} \int d^2x b_i(x) \cdot \hat{J}_i(x) &= \pi \bar{\rho} \epsilon_{ij} \left(\sum_{\alpha} q_{\alpha}^v \dot{x}_i^{v,\alpha} x_j^{v,\alpha} + \sum_{\alpha} q_{\alpha}^S \dot{x}_i^{S,\alpha} x_j^{S,\alpha} \right) \\ &\quad - \nu \left(\sum_{\alpha \neq \beta} q_{\alpha}^v q_{\beta}^v \epsilon_{ij} \dot{x}_i^{v,\alpha} \frac{(x^{v,\alpha} - x^{v,\beta})_j}{(x^{v,\alpha} - x^{v,\beta})^2} + \sum_{\alpha, \beta} q_{\alpha}^S q_{\beta}^v \epsilon_{ij} \dot{x}_i^{S,\alpha} \frac{(x^{S,\alpha} - x^{v,\beta})_j}{(x^{S,\alpha} - x^{v,\beta})^2} \right) \end{aligned}$$

$$+ \sum_{\alpha, \beta} q_{\alpha}^v q_{\beta}^S \epsilon_{ij} \dot{x}_i^{v, \alpha} \frac{(x^{v, \alpha} - x^{S, \beta})_j}{(x^{v, \alpha} - x^{S, \beta})^2} + \sum_{\alpha \neq \beta} q_{\alpha}^S q_{\beta}^S \epsilon_{ij} \dot{x}_i^{S, \alpha} \frac{(x^{S, \alpha} - x^{S, \beta})_j}{(x^{S, \alpha} - x^{S, \beta})^2} \Big). \quad (5.1)$$

It is obvious that the terms proportional to ν in (5.1) determine the statistics of solitons, because they are rewritten as⁵

$$\sum_{\alpha, \beta} \frac{d}{dt} \theta(x^{\alpha} - x^{\beta}), \quad (5.2)$$

where $\theta(x) = \arctan\left(\frac{x_2}{x_1}\right)$. In the path-integral formalism, this statistical term gives nontrivial contribution when world lines of two particles entwine each other. As we briefly mentioned in Sec.2 and Sec.3, similar terms appear from the coupling of solitons with the CS gauge field a_{μ}^{-} .

Anyonic properties of topological excitations plays an important role, for example, for hierarchical structure of the FQHE. Anyonic vortices and merons can be expressed in terms of boson fields with CS flux attached. At specific filling factors, these bosonized anyons can Bose condense, as the bosonized electrons. These states correspond to daughter states of the FQHE, and it is interesting to compare theoretical prediction of filling fractions of the FQHS with experimental observations[15].

In this section, we shall study quantum system of quasi-excitations in the FHQS. The first term of (5.1) determines canonical commutation relations (CCR) of soliton coordinates. Generally, the quantum system of anyons, whose Lagrangian is given by (5.1), is very difficult to solve. Therefore, we shall mostly concentrate on two-particle systems in the rest of discussion. In specific but physically interesting cases of two-particle systems, it can be shown that the statistical terms can be ignored in the first approximation, as we discuss later on. On the other hand, we shall take into account interactions between solitons.

Returnning to the real-time formalism, the Lagrangian of the multi-soliton system

⁵It should be remarked that in the present system, the size of solitons or the magnetic length itself plays a role of short-distance cutoff.

is given as,

$$\begin{aligned}\mathcal{L}_{\text{soliton}} = & -\pi\bar{\rho}\epsilon_{ij}\left(\sum_{\alpha}q_{\alpha}^v\dot{x}_i^{v,\alpha}x_j^{v,\alpha}+\sum_{\alpha}q_{\alpha}^S\dot{x}_i^{S,\alpha}x_j^{S,\alpha}\right) \\ & -\sum_{\alpha}\left(D(q_{\alpha}^v)+D(q_{\alpha}^S)\right)-\sum_{\omega,\omega'=v,S}\sum_{\alpha,\beta}(e\nu)^2q_{\alpha}^{\omega}q_{\beta}^{\omega'}W(x^{\omega,\alpha}-x^{\omega',\beta}),\end{aligned}\quad (5.3)$$

where $D(q_{\alpha}^v) = D_v$ and $D(q_{\alpha}^S) = D_m$ are energies to create the vortex and the meron, respectively (see (3.26)), and $W(x^{\omega,\alpha} - x^{\omega',\beta})$ is the Coulomb interactions between solitons. From (5.3), the CCR 's among soliton coordinates are given as[16],

$$\begin{aligned}[x_i^{v,\alpha}, x_j^{v,\beta}] &= -i\frac{\epsilon_{ij}}{2\pi\bar{\rho}q_{\alpha}^v}\delta_{\alpha\beta}, \\ [x_i^{v,\alpha}, x_j^{S,\beta}] &= 0, \\ [x_i^{S,\alpha}, x_j^{S,\beta}] &= -i\frac{\epsilon_{ij}}{2\pi\bar{\rho}q_{\alpha}^S}\delta_{\alpha\beta}.\end{aligned}\quad (5.4)$$

Here we have neglected the statistical terms. After obtaining quantum mechanical states of two-particle systems, we shall give plausible argument supporting this approximation. From (5.4), conjugate momentum of x_i is $\epsilon_{ij}x_j$. This obviously corresponds to the cyclotron motion in the classical picture. From this fact, dispersion relation of a pair of solitons is, in some cases, obtained quite easily.

Actually concerning vortices, the Lagrangian (5.3) has a similar structure with that of vortices in the single-layer system. It is rather straightforward to show that a vortex-antivortex pair, which are separated with each other more than the magnetic length, form a magneto-roton excitation. Its dispersion relation is given as[11]

$$E(p_t) = 2D_v - \frac{\nu^3 e^2}{p_t l_0^2}, \quad (5.5)$$

where p_t is the eigenvalue of the total momentum. This result is obtained by ignoring the statistical terms, but is in qualitative agreement with the results of the exact diagonalization[17].

Similar relation is obtained for a pair of merons. However, there is a very important effect which we have to take into account in the DL electron systems, i.e.,

inter-layer tunneling. Hamiltonian of the inter-layer tunneling is given as,

$$\begin{aligned}\mathcal{H}_T &= -\Delta_{\text{SAS}}(\bar{\psi}_\uparrow\psi_\downarrow + \bar{\psi}_\downarrow\psi_\uparrow) \\ &= -\Delta_{\text{SAS}}(\bar{\Psi}\sigma_1\Psi),\end{aligned}\tag{5.6}$$

where Δ_{SAS} is the energy splitting between symmetric and anti-symmetric states with respect to the pseudospin index.⁶

In terms of the parametrization (4.1),

$$\mathcal{H}_T = -\Delta_{\text{SAS}}J_0(\bar{z}\sigma_1z),\tag{5.7}$$

and therefore it is obvious that $\mathcal{H}_T$ gives an linear attractive confining force between $(-1, 0) - (0, -1)$ pair, $(-1, 0) - (1, 0)$ pair, etc., see Fig.5.

Let us first consider a pair of $(-1, 0)$ and $(1, 0)$ merons, whose coordinates are (x_1, y_1) and (x_2, y_2) , respectively. From (5.4), the nonvanishing CCR's are

$$[x_1, y_1] = iR, \quad [x_2, y_2] = -iR, \quad R \equiv \frac{2l_0^2}{\nu}.\tag{5.8}$$

Introducing the center of mass and relative coordinates as $X = \frac{1}{2}(x_1 + x_2)$, $Y = \frac{1}{2}(y_1 + y_2)$, $x = x_1 - x_2$ and $y = y_1 - y_2$, the CCR' are

$$[X, y] = iR, \quad [Y, x] = -iR.\tag{5.9}$$

From (5.9), it is obvious that $Y = R\hat{p}_x$ and $y = R\hat{p}_X$, where $\hat{p}_x$ and $\hat{p}_X$ are relative and total momentum operators, respectively. On the other hand, potential energy of this system is give as

$$V_1(r_{12}) = -\left(\frac{e\nu}{2}\right)^2 \frac{1}{r_{12}} + T_1 r_{12},\tag{5.10}$$

⁶Strictly speaking, the tunneling term (5.6) should be written in terms of the original fermionic electron operators. As a result, an extra factor like $\exp(2i\sigma_2 \int^x a_j^- dx_j)$ appears. This factor is important to preserve the gauge symmetry. However, results of the later discussion are not affected by this factor.

where $r_{12} = |\vec{r}_1 - \vec{r}_2|$ and tension $T_1 \propto \Delta_{\text{SAS}}$. If the total momentum of the pair is directed to the x -axis with magnitude $\hat{p}_X = p_X$, the energy is given as

$$E(p_X) = 2D_m - \left(\frac{e\nu}{2}\right)^2 \frac{\nu}{2l_0^2} \frac{1}{p_X} + \frac{2T_1 l_0^2}{\nu} p_X. \quad (5.11)$$

The above dispersion is correct for $r_{12} > l_0$ and $p_X > l_0/R$. Therefore, $(-1, 0) - (1, 0)$ meron pair also has magneto-roton dispersion. However, in contrast with the vortex-antivortex pair, its energy increase linearly with total momentum.

Let us turn to the $(-1, 0) - (0, -1)$ meron pair. Nontrivial CCR's are

$$[x_1, y_1] = iR, \quad [x_2, y_2] = iR. \quad (5.12)$$

From (5.12),

$$[X, Y] = \frac{i}{2}R, \quad [x, y] = 2iR. \quad (5.13)$$

In terms of the complex coordinates $\xi = x + iy$, $[\bar{\xi}, \xi] = -4R$, or by introducing radial and angle coordinates as $\xi = re^{i\varphi}$, the CCR is $[r^2, \varphi] = -4iR$. Energy of this system is given as

$$E(r) = 2D_m + \left(\frac{e\nu}{2}\right)^2 \frac{1}{r} + T_2 r, \quad (5.14)$$

where again $T_2 \propto \Delta_{\text{SAS}}$. It is obvious that merons are rotating around the center of mass with a fixed radius. The minimum of the energy is located at $r_m = \frac{e\nu}{2} \frac{1}{\sqrt{T_2}}$. Furthermore at long distance, meron pairs $(-1, 0) - (0, -1)$ and $(1, 0) - (0, 1)$ behave as vortices $(-1, -1)$ and $(1, 1)$, respectively.

There are classical pictures corresponding to the above eigenstates of $(-1, 0) - (1, 0)$ and $(-1, 0) - (0, -1)$ meron pairs. They are given in Fig.6. Merons are in the external magnetic field and also have electric charge propotional to their topological charge. Therefore, the Lorentz force works for them, as shown by the first and second terms in (5.3). This Lorentz force for merons is balanced by the Coulomb and the confining linear potentials between merons in a pair. Therefore, $(-1, 0) - (1, 0)$ meron pair moves with a constant velocity and a constant relative position. On the other hand in $(-1, 0) - (0, -1)$ pair, each meron rotates around the center of mass coordinate

with a constant cyclotron velocity. These classical pictures support the approximation which ignores the statistical terms. For $(-1, 0) - (1, 0)$ meron pair, the angle $\theta(\vec{r}_1 - \vec{r}_2)$ is a constant of motion in the classical limit and then $\frac{d}{dt}\theta(\vec{r}_1 - \vec{r}_2) = 0$. In the path-integral formalism, paths close to the classical path dominate. Furthermore, the statistical term is invariant under continuous deformations of path. Therefore for dominant paths, the statistical term does not contribute to the path integral in the present case. We think that this is the reason why the dispersion relation of vortex pair in the single-layer case given by (5.5) is, at least, qualitatively correct[17]. Similarly for $(-1, 0) - (0, -1)$ meron pair, $\theta(\vec{r}_1 - \vec{r}_2) \propto t$ in the classical picture, and the statistical term gives only irrelevant effects. Therefore quantum states of meron pairs which we obtained in the present discussion are consistent with the approximation which neglects the statistical terms. We hope that results of exact diagonalization of the double-layer systems support our results.

Here however it should be remarked that in the derivation of the Lagrangian of soliton (5.3), rotation of solitons etc. have not been taken into account, and therefore additional terms like $(\dot{x}_i^{\omega, \alpha})^2$, $\dot{x}_i^{\omega, \alpha} \ddot{x}_i^{\omega, \alpha}$, etc. must appear as higher-derivative terms in the effective theory of solitons, if we consider effects of them. For example, the additional term in the LLL condition $\epsilon_{ij} \partial_i \hat{J}_j$ induces terms like $(\dot{x}_i^{\omega, \alpha})^2$. This situation always happens in deriving an effective theory of topological objects with finite size, and most of cases low-energy behavior of these objects is well-described by the low-derivative model like (5.3), though many ideas have been proposed for modification of that. Especially for constant motion of $(-1, 0) - (1, 0)$ meron pair, higher-derivative terms can be legitimately ignored.

6 Conclusion

In this paper, we have discussed topological excitations in the CS gauge theory with complex boson fields, which describes DL FQHE. There are two types of topological

excitations in the LLL. The one is the vortices which have a nontrivial $U(1)$ winding number in the *common* phase factor of both pseudospin up and down electron fields. They are quite similar to the vortex of quasi-hole in the single-layer system. The others are the merons which are pseudospin textures and have a nontrivial $SU(2)$ winding number.

We first give qualitative arguments about these excitations and then numerical solutions. We derived an effective low-energy theory for them by using duality transformation and the LLL condition. It is shown that pseudospin textures are described by the CP^1 nonlinear σ -model with a CS gauge interaction and a Hopf term. The Hopf term determines statistical parameters of vortices and merons, and the result is consistent with that given by the argument through the Aharonov-Bohm effect. We also consider quantum system of multi-soliton configuration by retaining only center coordinates of solitons. Lagrangian of this quantum system is obtained from the effective low-energy Lagrangian of solitons. We took into account the effects of inter-layer tunneling, and determined energy spectrum of meron pairs. It is quite interesting to observe these excitations by experiment. Detailed study of dynamical properties of topological excitations is under study and results will be reported in future publications.

References

- [1] S.L.Sondhi, A.Karlhede, and S.A.Kivelson, Phys.Rev.**B47**(1993)16419;
H.A.Fertig, L.Brey, R.Cote, and A.H.MacDonald, Phys.Rev.**B50**(1994)11018.
- [2] S.E.Barrett, G.Dabbagh, L.N.Pfeiffer, K.W.West, and Z.Tycko,
Phys.Rev.Lett.**74**(1995)5112;
A.Schmeller, J.P.Eisenstein, L.N.Pfeiffer, and K.W.West,
Phys.Rev.Lett.**75**(1995)4290;
E.H.Aifer, B.B.Goldberg, and D.A.Broido, Phys.Rev.Lett.**76**(1996)680.
- [3] K.Moon, H.Mori, Kun Yang, S.M.Girvin, A.H.MacDonald, L.Zheng, D.Yoshioka,
and S-C.Zhang, Phys.Rev.**B51**(1995)5138;
Kun Yang, K.Moon, H.Mori, L.Belkhir, S.M.Girvin, A.H.MacDonald, L.Zheng
and D.Yoshioka, Phys.Rev.**B54**(1996)11644.
- [4] S.C.Zhang, H.Hansson, and S.Kivelson, Phys.Rev.Lett.**62**(1989)82;
N.Read, Phys.Rev.Lett.**62**(1989)86.
- [5] R.B Laughlin, Phys.Rev.Lett. **50**(1983)1395.
- [6] B.I.Halperin, Helv.Phys.Acta. **56**(1983)75.
- [7] Z.F.Ezawa and A.Iwazaki, Phys.Rev.**B47**(1993)7295.
- [8] Z.F.Ezawa, M.Hotta, and A.Iwazaki, Phys.Rev.**B46**(1992)7765.
- [9] Z.F.Ezawa, M.Hotta, and A.Iwazaki, Phys.Rev.**D44**(1991)452.
- [10] D-H.Lee and C.L.Kane, Phys.Rev.Lett.**64**(1990)1313.
- [11] M.P.A.Fisher and D.H.Lee, Phys.Rev.**B39**(1989)2756;
S.C.Zhang, Int.J.Mod.Phys.**B6**(1992)25.
- [12] F.Wilczek and A.Zee, Phys.Rev.Lett.**51**(1983)2250.

- [13] K.Kimm, K.Lee, and T.Lee, Phys.Rev.**D53**(1996)4436;
 Phys.Lett.**B380**(1996)303;
 K.Arthur, D.H.Tchrakian, and Yisong Yang, Phys.Rev.**54**(1996)5245.
- [14] I.Ichinose and A.Sekiguchi, paper in preparation.
- [15] I.Ichinose and A.Sekiguchi, paper in preparation.
- [16] A.L.Fetter, Phys.Rev.**162**(1967)143;
 F.D.M.Haldane and Y.-S.Wu, Phys.Rev.Lett.**55**(1985)2887.
- [17] F.D.M.Haldane, in “*The Quantum Hall Effect*” (Springer Verlag, 1990) editated
 by R.Prange and S.Girvin.

Figure Captions

Fig.1: Configurations of pseudospin vector of merons. Obviously, topological charges of them are fractional.


Fig.2: Numerical solutions of vortex, merons, etc., in $(3, 3, 2)$ state of the DL FQHE. In both upper and lower layers, charge densities have nontrivial behavior. This is a result of inter-layer Coulomb interaction.

Fig.3: Meron solution in the $(5, 5, 0)$ state. There are no correlations between upper and lower layers.


Fig.4: Vortex and merons in the $(1, 1, 4)$ state. Merons have a long undulating tail because of the dominant inter-layer Coulomb interaction.

Fig.5: Typical configuration of the $(-1, 0) - (0, -1)$ meron pair in the presence of the inter-layer tunneling. In the region between two merons, direction of pseudospins is reversed, and there is a confining linear potential between merons.

Fig.6: Corresponding classical pictures for systems of meron pairs. Lorentz force is balanced by the Coulomb and the linear interactions.




(a) $(-1,0)$ meron




(b) $(0,-1)$ meron


Figure 1: Configurations of pseudospin vector of merons. Obviously, topological charges of them are fractional.




(a) $(-1,-1)$ vortex



(b) $(-1,0)$ meron



(c) $(-2,0)$ meron



(d) $(-2,-1)$ meron

Figure 2: Numerical solutions of vortex, merons, etc., in $(3,3,2)$ state of the DL FQHE. In both upper and lower layers, charge densities have nontrivial behavior. This is a result of inter-layer Coulomb interaction.





Figure 3: Meron solution in the (5,5,0) state. There are no correlations between upper and lower layers.



(a) (-1,-1)vortex

(b) (-1,0)meron

Figure 4: Vortex and merons in the (1,1,4) state. Merons have a long undulating tail because of the dominant inter-layer Coulomb interaction.




Figure 5: Typical configuration of the $(-1,0)$ - $(0,-1)$ meron pair in the presence of the inter-layer tunneling. In the region between two merons, direction of pseudospins is reversed, and there is a confining linear potential between merons.






Figure 6: Corresponding classical pictures of the systems of meron pairs. Lorentz force is balanced by the Coulomb and the linear interaction.